

The Prevalence of Chlamydia Trachomatis Infection Among Gynecological Outpatients Attendees at Central Basra Hospitals Using One-Step Chlamydia Test

Ali Falih Al-Assadi*, Zaynab Yassin, Huda Salih Abood

Department of Obstetrics and Gynecology, Basra Medical College, Basra, Iraq

Correspondence to: Ali Falih Al-Assadi, Department of Obstetrics and Gynecology, Basra Medical College, Basra, Iraq; **E-mail:** alifalih1@gmail.com

Received date: April 10, 2020; **Accepted date:** April 18, 2020; **Published date:** April 28, 2020

Citation: Al-Assadi AF, Yassin Z, Abood HS (2020) The Prevalence of Chlamydia Trachomatis Infection Among Gynecological Outpatients Attendees at Central Basra Hospitals Using One-Step Chlamydia Test. J Obst Gynecol Surg. 1(1): pp. 1-6.

Copyright: ©2020 Al-Assadi AF, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

ABSTRACT

Objective: The objective of this study is to estimate the prevalence rate of chlamydia infection among gynecological outpatients attendees at central Basra hospitals and assessing the predisposing factors and clinical features.

Methods: This is a cross-sectional study that was conducted at central Basra hospitals during the period from 15 February 2018 to 10 May 2019. The distribution of cases according to the hospitals was 290 patients from Basra maternity and child hospital, 85 patients from Basra general hospital, 75 patients from AL Mawani hospital and 50 patients from AL Fayhaa hospital, this variation in the number depend on outpatient clinics attendees.

participants were assessed according to a predesigned questionnaire and screening test for Chlamydia Trachomatis was done for all patients under study by using one-step chlamydia test (Chlamydia Rapid Test Device).

Results: Among 500 patients 60% of them were from the age group 20-40 year, 62% were P2-4 and 89% from multipara and grand multipara, 60% were illiterate and 38% were having primary and secondary school, 61% from low socioeconomic class, nonusers of contraception were 23.7% and about 90% of patients were living at the central areas of Basra.

Several factors have a statistically significant effect on the Chlamydia Trachomatis infection like being infertile p-value 0.000 or having high parity p-value 0.013 or non-barrier contraceptives users p-value 0.035. While other factors have no statistically significant effect like age p-value 0.506, socioeconomic state p-value 0.779, and level of education p-value 0.986.

Conclusion: The prevalence rate of Chlamydia infection is low among the population in the central areas of Basra city compare to other countries. Being infertile or of high parity are among the significant risk factors. Barrier contraceptives significantly reduced the risk of infection and can be used to protect against the transmission of infection.

Keywords:

Chlamydia trochomatis, Gynecology, Basra Central Hospitals

Introduction

Chlamydia Trachomatis is the most common bacterial Sexually Transmitted Infections (STIs) worldwide, about 100 million cases of Chlamydia Trachomatis are detected annually worldwide according to the World Health Organization (WHO). In the united states alone, approximately 1.4 million Chlamydia infections occurred in 2013 [1,2]. Chlamydia is known as the "silent epidemic", as in women it may not cause any symptoms in 70-80% of cases and can remain for months or years before being discovered [3].

Residents based data from the united states, Australia, and the united kingdom guide that between 3 and 5% of people under 30 years of age will have a Chlamydia infection at any point in time [4-8]. Globally, in India, approximately 105.7 million cases for each year [9].

In other studies in Ethiopia, the prevalence rate for Chlamydia infection of the cervix was 5.9%. Among acquitted women joining the antenatal clinic in Benin City, Nigeria, a prevalence

rate of 13.3% was noted, while Nwanguma et.al reported a prevalence of 33% in asymptomatic volunteers in another Nigerian population [10-12]. The incidence of Chlamydia infections in women has augmented extremely from 79 to 467 per 100,000 between 1987 and 2003 [13].

Calculations of the incidence of disease consequence (specifically, PID, ectopic pregnancy, and infertility in women) are lacking, largely because there are very few natural-history studies of Chlamydia infections in humans. A methodical review trying to establish the attributable risk of infertility among women following genital Chlamydia infection concluded that there is formerly not sufficient evidence to accurately determine the population attributable risk [14].

Screening programs for Chlamydia have been set up to reduce transmission and reproductive tract morbidity in many developed countries. Recommended annual screening of all sexually active women aged 25 years or less by the United States Center for Disease Control (CDC) and prevention, as well as the screening of older women with risk factors (for example, those who have a new sex partner or multiple sex partners [15,16].

Age is the most common demographic associate of infection

with Chlamydia infection in women is young age (<20 years). Other factors associated with Chlamydia infection include unmarried status, black race, null parity, and poor socio-economic condition. Multiple sexual partners, a new sexual partner, a lack of use of barrier contraceptive devices and synchronized Gonococcal infection are also known to be related to Chlamydia infection [17,18].

The Screening tests include Chlamydia Trachomatis Nuclear Acid Amplification Test (NAAT) performed on an endocervical swab specimen that provides the highest sensitivity [19,20]. Chlamydia Kit test immune assay which was used in this study because of easy to be done, cheaper than the NAAT.

Because of ease of specimen management, improved test accuracy, suitability in specimen management, and ease of screening sexually active men and women, the NAAT has largely replaced culture, the historic gold standard for Chlamydia diagnosis, and the non-amplified probe tests [21].

Materials and Methods

This is a cross-sectional study that was conducted at Basra city during the period from 15 February 2018 to 5 May 2019. Patients have unselected attendance to the gynecological outpatient's clinics at the 4 main hospitals at Basra center (Basra maternity and child hospital, Basra general hospital, AL Mawani hospital, and AL-Fayhaa hospital) as these hospitals serve the majority of the population at Basra city. A total of 500 patients were

involved, 290 patient was from Basra maternity and child hospital as it is the major hospital in the city, 85 patient from Basra general hospital, 75 patients from AL-Mawani hospital, 50 patients from AL-Fyahaa hospital. Verbal consents were obtained from all participants.

A special questionnaire was designed for the study.

Patients attending the gynecological outpatient clinics with the following symptoms were included (dysuria, pelvic pain, acute pelvic pain), vaginal discharge, post-coital bleeding, intermenstrual bleeding, infertility).

Screening for chlamydia infection was done by using the Chlamydia kit test (Chlamydia Rapid Test Device) which was a rapid immunoassay test and manufactured by (Zhejiang, China 310030).

Analysis of the data was done using the Statistical Packages for Social Sciences (SPSS) version 24. Comparisons between variables and Chlamydia infection were performed by cross tab using the Chi-square test and fishers exact test. The bivariate Odds Ratio (OR) and chi-square test used to examine the association between variables and Chlamydia Trachomatis infection, a multiple logistic regression analysis was used to investigate the effect of predisposing factors on the Chlamydia infection, in all cases p-value <0.05 was considered to be significant.

Results

Table 1: Demographic characteristics of patients under study.

| Characteristics | Number | Percentage |
|--------------------------------|--------|------------|
| Age | | |
| <20 years | 111 | 22.20% |
| 20-40 years | 300 | 60% |
| >40 years | 89 | 17.80 % |
| Parity(among fertile patients) | | |
| P1 | 55 | 11% |
| P2-4 | 310 | 62% |
| >5 | 135 | 27% |
| Level of Education | | |
| Illiterate | 300 | 60% |
| Primary & secondary school | 191 | 38% |
| College & high education | 9 | 1.80% |
| Socioeconomic state | | |
| Low level | 305 | 61% |
| Middle level | 187 | 37% |
| High level | 8 | 1.60% |
| Address | | |
| Basra center | 453 | 90% |

According to Table 1, 60% of them were from the age group (20-40) year. 62% were (P2-4). 60% were illiterate and 38% were having primary and secondary school. 61% were from low

socioeconomic class and about 90% of patients were living at the central areas of Basra.

Table 2: Clinical features of patients under study.

| Symptoms and Signs | Number | Percentage |
|--------------------|--------|------------|
| Acute pelvic pain | 309 | 61.80% |

| | | |
|-------------------------|-----|--------|
| Post coital bleeding | 116 | 23.20% |
| Intermenstrual bleeding | 91 | 18.20% |
| Vaginal discharge | 226 | 45.20% |
| Infertility | 16 | 3.20% |
| Cervical erosion | 125 | 25% |
| Mucopurulent discharge | 29 | 5.80% |
| Pelvic tenderness | 76 | 15.20% |
| Multiple Complaints | 175 | 35% |

The commonest complaint was acute pelvic pain (61.8%), by multiple complaints (35%), cervical erosion (25%) and post-vaginal discharge being the second complaint (45.2%) followed coital bleeding (23.2%).

Table 3: Comparison of symptoms and signs distribution among patients with and without Chlamydia infection.

| Symptoms and signs | Chlamydia +ve | | Chlamydia -ve | | p-value |
|--------------------------|---------------|--------|---------------|--------|---------|
| | N | % | N | % | |
| Pelvic pain | 7 | 19.40% | 302 | 62% | 0.572 |
| Post coital bleeding | 5 | 13.80% | 111 | 22.80% | 0.316 |
| Inter menstrual bleeding | 1 | 2.70% | 90 | 18.50% | 0.479 |
| Vaginal discharge | 9 | 25% | 217 | 44.60% | 0.094 |
| Infertility | 2 | 5.50% | 14 | 87.50% | 0.061 |
| Cervical erosion | 5 | 13.80% | 120 | 24.60% | 0.327 |
| Mucopurulent Discharge | 2 | 5.50% | 27 | 5.50% | 0.171 |
| Pelvic tenderness | 5 | 13.80% | 71 | 14.60% | 0.034 |

The majority of patients with positive Chlamydia test had multiple complaints (76.90%), vaginal discharge was the commonest presenting symptom (69.2%), the second symptom was pelvic pain which was 53.8% while the commonest presentation of patients with negative Chlamydia test were infertility (87.5%)

and pelvic pain (26%). Although some clinical features were higher among cases with positive tests compared to cases with negative & vice versa with others but the differences between the 2 groups were statistically not significant.

Table 4: The relationship between Age and Chlamydia infection.

| Age | Chlamydia +ve | | Chlamydia -ve | | Total | Chi-Square | p-value |
|-------------|---------------|--------|---------------|--------|-------|------------|---------|
| | N | % | N | % | | | |
| <20 years | 0 | 0.00% | 111 | 22.70% | 111 | 6.16 | 0.046 |
| 20-40 years | 8 | 61.50% | 292 | 59.90% | 300 | | |
| >40 years | 5 | 38.40% | 84 | 17.20% | 89 | | |
| Total | 13 | | 487 | | | | |

The higher percentage of patients in both groups were among the age group (20-40) years. The percentage of patients with positive tests was higher than those with the negative test in age

groups 20-40 and 40 years, these differences were statistically not significant.

Table 5: The relationship between level of education and Chlamydia infection.

| Level of Education | Chlamydia +ve | | Chlamydia -ve | | Total | Chi-Square | p-value |
|------------------------------|---------------|-------|---------------|--------|-------|------------|---------|
| | N | % | N | % | | | |
| Illiterate | 10 | 0.769 | 290 | 59.50% | 300 | 1.68 | 0.432 |
| Primary and secondary school | 3 | 0.237 | 188 | 0.386 | 191 | | |
| College and High education | 0 | 0 | 9 | 0.018 | 9 | | |
| Total | 13 | | 487 | | 500 | | |

The majority of patients in both groups were from low socioeconomic class and the percentage of patients was higher in the test +ve group. While the percentages of patients with -ve test

was higher in both middle and higher socioeconomic class and these differences were statistically not significant.

Table 6: The relationship between Chlamydia infection and socio-economic state.

| Socio-economic State | Chlamydia +ve | | Chlamydia -ve | | Total | Chi-Square | p-value |
|----------------------|---------------|---------|---------------|---------|-------|------------|---------|
| | N | % | N | % | | | |
| Low level | 10 | 76.90 % | 295 | 60.50 % | 305 | 150.00 | 0.472 |
| Middle level | 3 | 23.70 % | 184 | 37.70 % | 187 | | |
| High level | 0 | 0% | 8 | 1.64 % | 8 | | |
| Total | 13 | | 487 | | | | |

The majority of patients in both groups were from low socio-economic class and the percentage of patients was higher in the test +ve group. While the percentages of patients with -ve

test was higher in both middle and higher socio-economic class and these differences were statistically not significant.

Table 7: The relationship between contraceptive usage and Chlamydia infection.

| Contraceptive users | | Chlamydia +ve | | Chlamydia -ve | | Total | Chi-Square | p-value |
|---------------------|-------------|---------------|--------|---------------|--------|-------|------------|---------|
| | | N | % | N | % | | | |
| Yes | Barrier | 0 | 0% | 102 | 20.90% | 102 | 565 | 0.016 |
| | Non-barrier | 10 | 5.30% | 177 | 36.30% | 187 | | |
| No | | 3 | 23.07% | 208 | 42.70% | 211 | | |
| Total | | 13 | | 487 | | | | |

Most of the infected patients were non-contraceptive users (23.07). Non-barrier method (5.30%) higher than the barrier

method (0%), these differences were statistically significant.

Table 8: The relationship between parity (fertile patients) and Chlamydia infection.

| Parity | Chlamydia +ve | | Chlamydia -ve | | Total | Chi-Square | p-value |
|----------|---------------|--------|---------------|--------|-------|------------|---------|
| | N | % | N | % | | | |
| P1 | 2 | 15.30% | 53 | 10.28% | 55 | 13.73 | 0.013 |
| P2-4 | 2 | 15.30% | 308 | 63.20% | 310 | | |
| P5 above | 9 | 69.20% | 126 | 25.80% | 135 | | |
| Total | 13 | | 487 | | | | |

The percentage of patients with +ve test was higher among those with P5 and more, while the percentage of patients with

-ve test was higher among those with P2-4. These differences were statistically highly significant.

Table 9: The relationship between address and Chlamydia infection.

| Address | Chlamydia +ve | | Chlamydia -ve | | Total | Chi-Square | p-value |
|-----------------|---------------|--------|---------------|--------|-------|------------|---------|
| | N | % | N | % | | | |
| Center of Basra | 10 | 76.90% | 443 | 90.90% | 453 | 2.93% | 0.087 |
| Rural area | 3 | 23.07% | 44 | 9.03% | 47 | | |
| Total | 13 | | 487 | | | | |

The majority of patients were from Basra Center but there was

no statistically significant effect of an address.

Table 10: Logistic Regression Analysis.

| Features | B | p-value | O.R | 95% C.I | |
|---------------------------------|-------|---------|------|---------|-------|
| | | | | Lower | Upper |
| Parity | 1.886 | 0.013 | 6.59 | 1.98 | 22.04 |
| Infertility | 4.229 | 0 | 6.8 | 4.7 | 7.93 |
| Non barrier Contraceptive users | 0.74 | 0.035 | 8.9 | 0.087 | 2.6 |

Discussion

Chlamydia trachomatis infections are the most commonly reported sexually transmitted bacterial infections in the USA, Europe and globally [22,23].

About 43 million new cases were detected in South-East Asia [24]. In this study, the hospital-based prevalence rate among symptomatic patients was 2.6%. In a study done at AL Ramadi

city, 3.75% of hospital attendance had new infection [25]. At AL Najaf city 58.2% of hospital attendance were infected [26]. In a study done at Bulgaria, 39.1% of symptomatic females had a positive test for Chlamydia Trachomatis [27], while in a tertiary care center in North India 19.9% of patients were infected [28]. These differences could be due to the differences in the sample sizes. Duration of study and the test used to detect the

infection. More than 50% of Chlamydia Trachomatis infected patients were asymptomatic [25]. All participants in this study were symptomatic because the study was hospital-based. The commonest complaints of infected patients was vaginal discharge (69.2%), followed by acute pelvic pain (53.8%), cervical erosion (38.5%), and post-coital bleeding (38.5%).

In women, the most common clinical features include abnormal vaginal discharge, vaginal bleeding (including bleeding after intercourse), and dysuria [29] chronic pelvic pain and pelvic tenderness (38.5%). While only 4% had chronic pelvic pain as reported by J.S. et al [23].

In this study, the percentages of all clinical features were higher in comparison to other studies [30-32] and this could be because all participants were having symptoms. Regarding the relationship between age group and Chlamydia infection, the majority (60%) of participants in this study among age groups from 20-40 years and this is similar to that reported by other studies done in Nigeria [33] and North India [28]. This is because those are the most sexually active and the infection is sexually transmitted. For the relationship between the level of education and infection, the highest prevalence of infected and non-infected patients were illiterate and it was higher among infected compared to non-infected patients. This could be due to their unawareness about the STD and the mode of transmission. In a study done in Nigeria, the highest prevalence was among university students [33].

The majority of infected patients in the current study were from the low socioeconomic class and this is similar to the finding of Okoror et al. [33]. The high-income earners can improve their health care and have access for early treatment of infection.

We reported a statistically significant relationship between contraceptive users and Chlamydia infection with 0% among barrier methods (protected) users. This is similar to that reported by Okoror et al. [33]. Patients of high parity (P5 and more) had a statistically highly significant higher rate of infection and this could be due to higher sexual activity. Because the study was conducted at Basra central hospitals there was no effect of address on the occurrence of infection and the majority of patients in both groups were from Basra central areas.

Conclusion

The prevalence rate of Chlamydia infection is low among the population in the central areas of Basra city compared to other countries. Being infertile or of high parity are among the significant risk factors.

Barrier contraceptive (Condom) significantly reduces the risk of infection and can be used to protect against the transmission of infection.

References

1. World Health Organization (2011) Incidence of selected sexually transmitted infections. Chlamydia trachomatis, Neisseria gonorrhoeae, syphilis and Trichomonas vaginalis: methods and results used by WHO to generate 2005 estimates. W. Press. Editor.
2. Panelists GG, Taylor M, Berman B, et al. (2016) Sinecatechins ointment. 15% for the treatment of external genital and

perianal warts: proceedings of an expert panel roundtable meeting. *J Clin Aesthet Dermatol.* 9(3): pp. S2-S15.

3. NHS over view chlamydia Page last reviewed: 4 June 2018 .
4. Lewis D, Newton DC, Guy RJ, et al. (2012) The prevalence of Chlamydia trachomatis infection in Australia: a systematic review and meta-analysis. *BMC infectious diseases.* 12(1): pp. 113.
5. Sonnenberg P, Clifton S, Beddows S, et al. (2013) Prevalence, risk factors, and uptake of interventions for sexually transmitted infections in Britain: Findings from the National Surveys of Sexual Attitudes and Lifestyles (Natsal). *The Lancet.* 382(9907): pp. 1795-806.
6. Datta SD, Sternberg M, Johnson RE, et al. (2007) Gonorrhoea and chlamydia in the United States among persons 14 to 39 years of age 1999 to 2002. *Ann Intern Med* 147(2): pp. 89-96.
7. Datta SD, Torrone E, Kruszon-Moran D, et al. (2012) Chlamydia trachomatis trend in the United States among persons 14 to 39 years of age 1999-2008. *Sexually transmitted diseases.* 39(2): pp. 92-6.
8. Price MJ, Ades AE, De Angelis D, et al. (2014) Incidence of Chlamydia trachomatis infection in women in England: two methods of estimation. *Epidemiol Infect.* 142(3): pp. 562-576.
9. World Health Organization (2012) Global incidence and prevalence of selected curable sexually transmitted infections-2008.
10. Buve A, Weiss HA, Laga M, et al. (2001) The epidemiology of gonorrhoea, chlamydial infection and syphilis in four African cities. *Aids.* 15(1):S79-88.
11. Isibor JO, Ugbomoiko D, Nwobu GO, et al. (2005) Detection of chlamydia antigen in cervical specimens from antenatal clinic attendees in Benin City, Nigeria. *African J Clin Exp Microbiol.* 6(3): pp. 208-211.
12. Nwanguma BC, Kalu I, Ezeanyika LU (2009) Seroprevalence of Anti-chlamydia trachomatis IgA antibody in a Nigerian population: Diagnostic significance and implications for the heterosexual transmission of HIV. *Int J Infect Dis.* 7(2): pp. 8-15.
13. Malhotra M, Sood S, Mukherjee A, et al. (2013) Genital Chlamydia trachomatis: an update. *The Indian journal of medical research.* 138(3): pp. 303.
14. Wallace LA, Scoular A, Hart G, et al. (2008) What is the excess risk of infertility in women after genital chlamydia infection? A systematic review of the evidence. *Sexually Transmitted Infections.* 84(3): pp. 171-175.
15. Oakshott P, Kerry S, Aghaizu A. et al. (2010) Randomised controlled trial of screening for Chlamydia trachomatis to prevent pelvic inflammatory disease: the POPI (prevention of pelvic infection) trial. *Bmj.* 8(1): pp. c1642.
16. Workowski KA, Berman SM, Douglas JM (2008) Emerging antimicrobial resistance in Neisseria gonorrhoeae: urgent need to strengthen prevention strategies. *Annals of internal medicine.* 148(8): pp. 606-613.
17. Novak M, Novak D (2013) Risk factors for Chlamydia trachomatis infection among users of an internet-based testing service in Sweden. *BMJ Sex Reprod Health.* 4(1): pp. 23-27.
18. Batteiger BE (2012) Chlamydia infection and epidemiology. *In Intracellular Pathogens I: Chlamydiales.* Am Soci Microb. pp. 1-26.

19. Black CM, Marrazzo J, Johnson RE, et al. (2002) Head-to-head multicenter comparison of DNA probe and nucleic acid amplification tests for Chlamydia trachomatis infection in women performed with an improved reference standard. *J Clin Microb.* 40(10): pp. 3757-3763.
20. Sary A (1999) Correct samples for diagnostic tests in sexually transmitted diseases: which sample for which test?. *FEMS Immunol Med Mic.* 24(4): pp. 455-459.
21. Centers for Disease Control (1993) Recommendations for the Prevention and Management of Chlamydia Trachomatis Infections 1993. *MMWR.* 42(RR1-2): pp. 1-37.
22. Newman L, Rowley J, Vander Hoorn S (2015) Wivesooriya NS, Unemo M. infections in 2012 based on systematic review and global reporting. *PLoS one.* 10(12): pp. e0143304.
23. Wilson JS, Honey E, Templeton A, et al. (2002) A systematic review of the prevalence of Chlamydia trachomatis among European women. *Hum Reprod.* 8(4): pp. 385-394.
24. World Health Organization (2001) Global prevalence and incidence of selected curable sexually transmitted infections: overview and estimates. Geneva: World Health Organization.
25. Al-Alouci MM, Al-Hayani NN, Al-Alouci TM (2019) Assessment of Chlamydia trachomatis infection in symptomatic women by ELISA and evaluate the levels of CRP, C3, C4 and IgA in patients sera. *J Pharm Sci & Res.* 11(3): pp. 1131-1135.
26. Al-Abbas WD, Radhi OA (2019) Incidence of Chlamydia trachomatis and Trichomonas Vaginalis Genital Infections among Non-Pregnant Women in Al-Najaf Province. *Kufa J Nurs sci.* 9(1): pp. 1-8
27. Wilson JS, Honey E, Templeton A, et al. (2002) Sary A, Stray-Pedersen B. A systematic review of the prevalence of Chlamydia trachomatis among European women. *HumReprod.* 8(4): pp. 385-394.
28. Malhotra M, Bala M, Muralidhar S (2008) Prevalence of Chlamydia trachomatis and its association with other sexual transmitted infections in a tertiary center in North India. STD Teaching Training and Research center. Vardhman Mahavir medical college, New Delhi, India.
29. Stamm WE (1999) Chlamydia trachomatis infections of the adult. *Sexually transmitted diseases.*
30. Gonzales GF, Munoz G, Sanchez R, et al. (2004) Update on the impact of Chlamydia trachomatis infection on male fertility. *Andrologia.* 36(1): pp. 1-2.
31. Hu D, Hook EW, Goldie SJ (2004) Screening for Chlamydia trachomatis in women 15 to 29 years of age: a cost-effectiveness analysis. *Ann Int Medic.* 141(7): pp. 501-513.
32. Schneede P, Tenke P (2003) Hofstetter AG. Sexually transmitted diseases (STDs)-A synoptic overview for urologists. *European Urology.* 44(1): pp. 1-7.
33. Ehis OL, Cyril O, Tolulope E, et al. (2014) Prevalence and risk of Chlamydia trachomatis in symptomatic patients attending clinics in South West Nigeria. *J Clin Microbiol.* 5(1): pp.1-1.